



Test Report

Report Number: SSMT-R-2022-04967-01B

Sample Name: eResin-PLA Pro

Study Title: Skin Irritation Test - 0.9% Sodium Chloride

Injection Extract

Standard: ISO 10993-10:2010

Test facility

Jiangsu Science Standard Medical Testing Co., Ltd.

C4 Building, No.9 Changyang Road, Wujin District, Changzhou, Jiangsu, China

Sponsor

Shenzhen Esun Industrial Co., Ltd.

Wuhan University Building A403-I and A901, No.6 Yuexing 2 Road, Nanshan District, Shenzhen, China

Jiangsu Science Standard Medical Testing Co., Ltd.

C4 Building No.9 Changyang Road, Wujin District, Changzhou, Jiangsu, China 213161 Tel: (86-519-83587899) Fax: (86-519-83587899) www.jcssmt.com Document No.: SHT-ASS-A11 Version 2.0 Page 1 of 11

Soline Science

Contents

Explanation	3
Conclusion	4
Study verification and signature	5
1.0 Purpose	6
2.0 Reference	6
3.0 Test and control articles	6
4.0 Identification of test system	7
5.0 Animal Care and Maintenance	7
6.0 Justification of the test system	7
7.0 Instruments and pH test paper	8
8.0 Experiment design and dose	8
9.0 Evaluation criteria	10
10.0 Results of the test	10
11.0 Deviation statement	10
12.0 Record	10
13.0 Confidentiality agreement	11

Explanation

- 1. Please apply for rechecking within 15 days of receiving the report if there is any objection.
- 2. Any erasure or without special testing seal renders the report null and void.
- 3. The report is only valid when signed by the persons who edited, checked and approved it.
- 4. The result relate only to the articles tested.
- 5. The report shall not be reproduced except in full without the written approval of the institute.
- 6. This experiment was carried out in the sub-site and the address is: E3 Building, No.9 Changyang Road, Wujin District, Changzhou, Jiangsu, China.

Document No.: SHT-ASS-A11 Version 2.0 Page 3 of 11

Conclusion

The animal skin irritation test was conducted to assess the potential irritation of the test article or material.

The test sample was extracted with 0.9% sodium chloride injection. The patches (about 2.5 cm×2.5 cm) which moistened by 0.5 ml extract of test article were directly applied to the rabbit skin for 4 hours. Observation for erythema and edema were conducted at 1 h, 24 h, 48 h and 72 h after removal of the patches.

The skin reaction on test sites did not exceed that on the control sites. The primary irritation index for the test article was calculated to be 0.

The test result showed that the extract of the test article did not induce skin irritation in rabbit under the test condition.

Study verification and signature

The study was carried out in accordance with the standard operating procedure. The test process was conducted in compliance with the requirements of CNAS-CL01:2018 (ISO/IEC17025:2017, IDT) and RB/T214-2017.

Date Received	2022-08-24
Technical Initiation Date	2022-09-03
Technical Completion Date	2022-09-09
Final Report Completion Date	2022-09-13

Edited by	Churshay Yay	2022.09.13
		Date
Checked by	Suri Han	2022.09.13
		Date
Approved by	Dairy They Authorized signatory	2022.10.17
	Authorized signatory	Date

Jiangsu Science Standard Medical Testing Co., Ltd.

1名に とから

1.0 Purpose

New Zealand white rabbits were used to evaluate the potential of skin irritation of samples under the conditions of this test.

2.0 Reference

Biological evaluation of medical devices-Part 10:Tests for irritation and skin sensitization(ISO 10993-10:2010)

3.0 Test and control articles

3.1 Test article (The information about the test article was supplied by the sponsor wherever applicable.)

Test article name: eResin-PLA Pro

Initial State: Non-sterile

Model: N/S Size: N/S

Lot/ Batch#: N/S

Physical State: Solid Color: See the photo

Density: N/S Stability: N/S Solubility: N/S

Test Article Material: N/S Packing Material: N/S

Storage Condition: Room temperature

Manufacturer: Shenzhen eSunMed Biotechnology Co.,Ltd.

Manufacturer Address: Floor 3, No. 9, Yifenghua Innovation Industrial Park, Xinshi community, Dalang Street,

Longhua district, Shenzhen City

Sample photograph:



3.2 Control Articles

Name: 0.9% sodium chloride injection (SC)

Manufacturer: Anhui Shuanghe Pharmaceutical Co., Ltd.

Size: 500 ml

(C) =

Physical State: Liquid

Color: Colourless

Lot/ Batch#: 200714 2C

Storage Condition: Room Temperature

4.0 Identification of test system

Species: New Zealand white rabbit

Number: 3 Sex: Female

Weight: Initial body weight not less than 2.0 kg

Health status: Healthy, young adult, nulliparous and not pregnant.

Housing: Animals were housed in groups in cages identified by a card indicating the lab number and test

code.

Animal identification: Cage card The quarantine period: 3 days

5.0 Animal Care and Maintenance

Animal purchase: Provided by Danyang Changyi experimental animal breeding Co., Ltd <Permit Code: SCXK (SU) 2021-0002>

Bedding: NA

Feed: Rabbit Diet, Beijing Keao Xieli Feed Co., Ltd.

Water: Drinking water met the Standards for Drinking Water Quality (GB 5749-2006)

Cages: Stainless steel cage, Suzhou Fengqiao purification equipment Co., Ltd.

Environment: Temperature 16-26°C, Relative humidity 40%-70%, Lights 12 hours light/dark cycle

Personnel: Associates involved were appropriately qualified and trained

Selection: Only healthy animals were selected

Veterinarian: Vet takes care of the whole course

Ethics: Test methods of operation were reviewed and approved by the Commission on Science Standard animal ethics

There were no known contaminants present in the feed, water expected to interfere with the test data.

6.0 Justification of the test system

6.1 The rabbit is specified as an appropriate animal model for evaluating potential skin irritants by the current testing standards. Positive control 15 % sodium dodecyl sulfate has been substantiated at SSMT with this method. Positive control test is conducted every six months. The last irritation index was 4.4(polar test group). The data was from the report SSMT-R-2022-02748-01(Complete Date: 2022.06.03).

6.2 The test article extract was directly applied to the rabbit skin, which is considered to be the best mean of contact.

Document No.: SHT-ASS-A11 Version 2.0 Page 7 of 11

7.0 Instruments and pH test paper

7.1 Instruments

Thermostatic oscillation incubator(SSMT-564)

Electronic balance (SSMT-075)

Clean bench (SSMT-501)

Electronic balance (SSMT-147)

7.2 PH test paper

Supplier: China Shanghai Sanai Si Reagent Co., Ltd

Lot/ Batch#: 20211103

8.0 Experiment design and dose

8.1 Sample preparation

The test article was extracted as Table 1. Extract was checked and used immediately after extraction without the process of filtering, centrifugation, dilution, etc. The pH of the extract was not adjusted prior to testing. The preparation process was aseptic. The control article was prepared under the same condition.

Table 1 Sample Preparation

Aseptic Sam	Aseptic Sampling Aseptic Agitation Extraction In Inert Container			Final Extract			
Sampling Manner	Actually Sampling	Extraction solvent	Extraction ratio	Solvent volume	Condition	Clear or Not	pН
Random sampling	3.65 g	0.9% sodium chloride injection	0.2 g : 1 ml	18.2 ml	37 ℃, 72 h	Clear	6.5~7.0


8.2 Test method

Use the rabbits with healthy intact skin. Fur is generally clipped on the back of the rabbits 16 h before testing, a sufficient distance on both sides of the spine for application and observation of all test sites (approximately 10×15 cm).

Apply 0.5 ml extract of test article or control to $2.5 \text{ cm} \times 2.5 \text{ cm}$ absorbent gauze patches, and then apply the patch soaked with the extract of test article or control directly to the skin on each side as shown in Figure 1. And then wrap the application site with a bandage (semi-occlusive) for 4 h. At the end of the contact time, remove the dressings.

Document No.: SHT-ASS-A11 Version 2.0 Page 8 of 11





1- Cranial end, 2- Test site, 3- Control site, 4- Clipped dorsal region, 5- Caudal end Figure 1 Location of skin application sites

8.3 Observation of animal

Describe and score the skin reaction for erythema and oedema according to the scoring system given in Table 2 for each application site at each time interval. Record the appearance of each application site at 1 h, 24 h, 48 h and 72 h following removal of the patches.

Table 2 Classification System for Skin Reaction

Reaction	Irritation score	
Erythema and Eschar Formation		
No erythema	0	
Very slight erythema (barely perceptible)	1	
Well-defined erythema	2	
Moderate erythema	3	
Severe erythema (beet redness) to eschar formation preventing grading of erythema	4	
Oedema Formation	7/X	
No oedema	0	
Very slight oedema (barely perceptible)	1	
Well-defined oedema (edges of area well-defined by definite raising)	2	
Moderate oedema (raised approximately 1mm)	3	
Severe edema (raised more than 1mm and extending beyond exposure area)	4	
Maximal possible score for irritation	8	

NOTE: Other adverse changes at the skin sites were recorded and are reported.

8.4 Result calculation

Use only 24 h, 48 h and 72 h observations for calculation.

After the 72 h grading, all erythema grades plus oedema grades 24 h, 48 h and 72 h are totalled separately for each test sample and blank for each animal. The primary irritation score for an animal is calculated by dividing the sum of all the scores by 6 (two test/observation sites, three time points).

To obtain the primary irritation index for the test sample add all the primary irritation scores of the individual animals and divide by the number of animals.

Calculate the primary irritation score for the controls and subtract that score from the score using the test material to obtain the primary irritation score.

9.0 Evaluation criteria

The primary irritation index is characterized by number (score) and description (reaction category) given in Table 3.

Table 3 Primary irritation index categories in a rabbit

Mean score	reaction category
0-0.4	Negligible
0.5-1.9	Slight
2.0-4.9	Moderate
5-8	Severe

10.0 Results of the test

According to what observed, the reaction of skin on testing side did not exceed that on the control side. Thus, the primary irritation index for the test article was calculated to be 0. See Table 4.

Table 4 Dermal observations

Rabbit No.	iroup	Interval				
Navolt No.		iroup	1h	24h	48h	72h
	Test	Erythema	0	0	0	0
01-J1501 Negati	Article	Oedema	0	0	0	0
	Negative	Erythema	0	0	0	0
	Control	Oedema	0	0	0	0
	Test	Erythema	0	0	0	0
2022-04967-	Article	Oedema	0	0	0	0
01-J1502 Negative Control	Erythema	0	0	0	0	
	Control	Oedema	0	0	0	0
Test 2022-04967- Article 01-J1503 Negative Control	Test	Erythema	0	0	0	0
	Article	Oedema	0	0	0	0
	Negative	Erythema	0	0	0	0
	Control	Oedema	0	0	0	0

Under the conditions of this study, the extract of the test article did not induce skin irritation in rabbit skin.

11.0 Deviation statement

There was no deviation from the standard operating procedure which were judged to have any impact on the validity of the data.

12.0 Record

All the original data and records related to this test and copies of the final report are retained in the archives of Science Standard Medical Testing.

Document No.: SHT-ASS-A11 Version 2.0

Page 10 of 11

13.0 Confidentiality agreement

Statements of confidentiality were as agreed upon prior to study initiation.



Document No.: SHT-ASS-A11 Version 2.0 Page 11 of 11